



# The Future of Emergency Medicine as Revealed through Research



*A Window into  
the Near Future*

PHIC CONTROLS CORPORATION

BUFFALO, NEW YORK

# **Raymond L. Fowler, M.D., FACEP**

**Co-Principal Investigator  
National Institutes of Health  
Resuscitation Outcomes Consortium**

**Joint Investigator  
National Heart, Lung, and Blood Institute  
IMMEDIATE Trial**

**Deputy Medical Director  
Dallas Area BioTel System**

**Co-Chief in the Section on  
EMS, Disaster Medicine, and Homeland Security  
Southwestern Medical Center**



A photograph of several parachutes descending against a sunset sky. The sky transitions from a deep orange at the bottom to a lighter yellow at the top. The parachutes are dark against the bright background. The entire image is framed by a thick dark blue border.

[www.uts.w.w.s](http://www.uts.w.w.s)

[www.rayfowler.com](http://www.rayfowler.com)

# My Perspective

*Save the whales: Collect the whole set!*

*42.7% of all statistics  
are made up on the spot*

*99% of lawyers  
give the rest a bad name*

*I intend to live forever....  
.....so far, so good*





**CNN.com** WITH FREE VIDEO Member Services Internet

SEARCH  THE WEB  CNN.com  SEARCH

Home Page  
**World**  
U.S.  
Weather  
Business at CNN Money  
Sports at Sports Illustrated  
Politics  
Law  
Technology  
Science & Space  
Health  
Entertainment  
Travel  
Education  
Special Reports  
Video **NOW FREE**  
Autos with Edmunds.com

**WORLD**

**London Terror** SPECIAL REPORT

[» Victims](#) | [Bombing map](#) | [Video log](#) | [Your e-mails](#) | [Eyewitness](#)

## Atta's father praises London bombs

Wednesday, July 20, 2005; Posted: 8:57 a.m. EDT (12:57 GMT)

**CAIRO, Egypt (CNN) --** The father of one of the hijackers who commandeered the first plane that crashed into the World Trade Center on September 11, 2001, praised the recent terror attacks in London and said many more would follow.

Speaking to CNN producer Ayman Mohyeldin Tuesday in his apartment in the upper-middle-class Cairo suburb of Giza, Mohamed el-Amir said he would like to see more attacks like the July 7 bombings of three London subway trains and a bus that killed 52 people, plus the four bombers.




**Mohamed Atta is believed to have piloted one of the planes that hit the World Trade Center.**

advertiser links: [what's this?](#)

SERVICES



## MADRID TRAIN BOMBINGS





# WORLD

Tools: [Save](#) | [Print](#) | [E-mail](#) | [Most Popular](#)

## Overloaded bus plunges off cliff, dozens killed

POSTED: 11:08 a.m. EDT, October 28, 2006

### ADVERTISER LINKS

- [Distance Learning](#)
- [Get a New Car Cheap](#)
- [Women's Apparel](#)

#### [Calculate Your New Payment](#)

\$510,000 mortgage for \$1,639/mo. Think you pay too much? Refi now.  
[www.lowermybills.com](http://www.lowermybills.com)

#### [Protiviti Risk Management Consulting](#)

Protiviti provides independent business risk, technology risk, and internal...  
[protiviti.com](http://protiviti.com)

#### [Become a Child Crisis Partner](#)

Your \$20 pledge helps one child after another escape lives of horror.  
[www.worldvision.org](http://www.worldvision.org)

[what's this?](#)

Adjust font size:

**KATMANDU, Nepal (AP)** -- An overcrowded bus plunged off a mountain road in western Nepal on Saturday, leaving at least 42 people dead and 45 injured, police said.

The bus was traveling near the town of Tribeni, some 250 miles west of Katmandu, when it skidded off the road, plunging about 950 feet into a ravine below, police officer Birkha Bahadur Rawal said.

Police were investigating the cause, but Rawal said an initial probe indicated the driver may have lost control of the vehicle because it was overloaded with passengers. It was not known whether the driver survived, police said.

Some of the passengers jumped off the bus before it crashed and survived, indicating several people were riding on the roof, a common practice in rural Nepal where not much public transportation is available.

A ticket taker on the bus, Jeevan Nepali, said from his hospital bed that the bus was unable to pick up speed on the slopes and the driver stopped. The bus then rolled backward and veered off the road into the ravine.

The bus was traveling toward the southern town of Dang, when the crash occurred.

W  
PAR  
CUR  
AUTO  
HAV  
L  
HAS

# DISASTER



## CNN.com

CNN.com

close window

### NORTH KOREA MISSILE RANGE

How far could North Korea launch a nuclear warhead? North Korea's longest-range missile, the Taepodong-2, has an estimated reach from 3,750 kilometers to 15,000 kilometers (2,325 miles to 9,300 miles). When test fired in July, a Taepodong-2 failed about 40 seconds after launch, landing in the Sea of Japan. North Korea's missile arsenal also includes the Nodong, with an estimated range of 1,300 kilometers (800 miles), and the Scud-C, with a range of 300 kilometers (185 miles).




[VIEW 1](#) [VIEW 2](#)

◀ Previous Next ▶

The apparent nuclear test was conducted at 10:36 a.m. (1:36 a.m. GMT) in

*perpetuating an arms race?*





The emerging of  
a subspecialty:



# Approaching the Patient





**“See what you see!”**



**“People look, but they  
don’t see”**

**...A. Fowler, Jr.**

**Alertness?**

**Level of distress?**

**Noises?**

**Respirations?**

**The pulse rate?**


**Skin?**

**Obvious things (bleeding)**





**Our pulse can only go  
so fast under sympathetic  
stimulation:**




**220 minus age**

$$\text{Baby} = (220 - 0) = 220$$


$$\text{Snerd} = (220 - 53) = 167$$

$$\text{Aunt Minnie} = (220 - 70) = 150$$






# What is this rhythm?



**Correct answer:**

**"This HAS to be  
an arrhythmia**

# What is this rhythm?



**Correct answer:**

**“It COULD be sinus tach”**



**If you forget everything  
else that I say:**

**Remember that  
patients having  
near maximum  
sinus tachycardia  
at rest  
are dying!**

**A “physiological  
response”**



***Something  
mobilizing a  
massive  
physiological  
response***

**Your job is  
to determine if  
a rapid rhythm  
MAY be sinus tach**

***If it is,  
you must take action***

**Because so many  
courses are too long,  
too boring,  
and teach difficult concepts  
to providers  
who will rarely use  
that information**





- **Airway**
- **Breathing**
- **Circulation**
  - **Drugs**
  - **Disaster**
- **Electrocardiography**

**Airway**



**It is not at all clear  
what the best airway devices  
are now or  
what they will be**

## **Paramedic Drug Assisted Intubation (DAI) in Georgia**

### Overview

The attached annotated bibliography contains most of the significant literature covering prehospital intubation and prehospital drug assisted intubation (DAI). Review of this literature will help to develop a policy on prehospital airway management and prehospital DAI.

### Fundamental Questions

- What is the definition of efficacious in the context of prehospital ETI?
- Is prehospital ETI feasible or efficacious?
- Is prehospital ETI facilitated with medications safe and efficacious?
- Can the characteristics of a safe and efficacious prehospital medication facilitated ETI program be defined and quantified?

Davis, D. P., J. Peay, et al. (2005). "The impact of prehospital endotracheal intubation on outcome in moderate to severe traumatic brain injury." Journal of Trauma-Injury Infection & Critical Care 58(5): 933-9.

**CONCLUSION:**

Prehospital intubation is associated with a decrease in survival among patients with moderate-to-severe TBI. More critically injured patients may benefit from prehospital intubation but may be difficult to identify prospectively.



Davis, D. P., J. Stern, et al. (2005). "A follow-up analysis of factors associated with head-injury mortality after paramedic rapid sequence intubation." Journal of Trauma-Injury Infection & Critical Care 59(2): 486-90.

**CONCLUSION:**

Paramedic RSI was associated with an increase in mortality compared with matched historical controls. The association between hyperventilation and mortality was confirmed. In addition, patients transported by helicopter after paramedic RSI had improved outcomes. Paramedic RSI did not seem to prevent aspiration pneumonia.

Mort, T. C. (2004). "Emergency tracheal intubation: complications associated with repeated laryngoscopic attempts.[see comment]." Anesthesia & Analgesia 99(2): 607-13.

intubate the trachea outside the operating room. There was a significant increase in the rate of airway-related complications as the number of laryngoscopic attempts increased ( $\leq 2$  versus  $> 2$  attempts): hypoxemia (11.8% versus 70%), regurgitation of gastric contents (1.9% versus 22%), aspiration of gastric contents (0.8% versus 13%) bradycardia (1.6% versus 21%), and cardiac arrest (0.7% versus 11%,  $P < 0.001$ ). Although predictable, this analysis provides data that confirm the number of laryngoscopic attempts is associated with the incidence of airway and hemodynamic adverse events. These data support the recommendation of the ASA Task Force on the Management of the Difficult Airway to limit laryngoscopic attempts to three in lieu of the considerable patient injury that may occur.

Silvestri, S., G. A. Ralls, et al. (2005). "The effectiveness of out-of-hospital use of continuous end-tidal carbon dioxide monitoring on the rate of unrecognized misplaced intubation within a regional emergency medical services system." Annals of Emergency Medicine 45(5): 497-503.

**CONCLUSION:** No unrecognized misplaced intubations were found in patients for whom paramedics used continuous ETCO<sub>2</sub> monitoring. Failure to use continuous ETCO<sub>2</sub> monitoring was associated with a 23% unrecognized misplaced intubation rate.



Ufberg, J. W., J. S. Bushra, et al. (2005). "Aspiration of gastric contents: association with prehospital intubation." American Journal of Emergency Medicine 23(3): 379-82.

9.08). Patients endotracheally intubated in the PH setting are more likely to have aspirated gastric contents than those intubated in the ED.

# **POSITION PAPER**

NATIONAL ASSOCIATION OF EMS PHYSICIANS

## **RECOMMENDED GUIDELINES FOR UNIFORM REPORTING OF DATA FROM OUT-OF-HOSPITAL AIRWAY MANAGEMENT:**

**POSITION STATEMENT OF THE NATIONAL ASSOCIATION OF EMS PHYSICIANS**

Henry E. Wang, MD, MPH, Robert M. Domeier, MD, Douglas F. Kupas, MD,  
Mark J. Greenwood, DO, JD, Robert E. O'Connor, MD, MPH

Henry E. Wang, MD, MPH, Robert M. Domeier, MD, Douglas F. Kupas, MD,  
Mark J. Greenwood, DO, JD, Robert E. O'Connor, MD, MPH

Dr. Wang is at the Department of Emergency Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania; Dr. Domeier is at the Department of Emergency Medicine, St. Joseph Mercy Hospital, Ann Arbor, Michigan; Dr. Kupas is at the Department of Emergency Medicine, Geisinger Health System, Danville, Pennsylvania; Dr. Greenwood is at the Department of Medicine, Section of Emergency Medicine, University of Chicago, Chicago, Illinois, and AeroMed at Spectrum Health, Grand Rapids, Michigan; and Dr. O'Connor is at the Department of Emergency Medicine, Christiana Care Health System, Newark, Delaware.

## CONCLUSION

Airway management, including endotracheal intubation, is the most important procedure performed in the prehospital setting. EMS services should closely monitor the performance of ETI to ensure that the highest level of care is provided. EMS services should adhere to the recommended standards for defining, collecting, and reporting airway management data. Although there are many methods for collecting airway management data, systems should use methods that result in the most accurate reports of treatment courses and outcomes.

## RECOMMENDED FORMAT FOR REPORTING SYSTEM-WIDE PERFORMANCE OF AIRWAY MANAGEMENT

Prehospital services should use the following guidelines for summarizing systemwide performance of airway management:

1. ETI success rate (percentage and relative frequency) for all ETI (pooled [based on overall outcome of patient encounter], not per attempt)
2. ETI success rate (percentage and relative frequency) for subset of patients in cardiac arrest
3. ETI success rate (percentage and relative frequency) for subset of patients with a pulse (nonarrest)
4. For patients with a pulse (non-arrest), ETI success rates (percentage and relative frequency) stratified by overall ETI method:
  - a. Orotracheal
  - b. Nasotracheal
  - c. Sedation-facilitated intubation
  - d. Rapid-sequence intubation
5. ETI success rates (percentage and relative frequency) for subset of pediatric patients (<18 years of age) (Individual services may choose to further stratify this group by specific age ranges.)
6. ETI success rates (percentage and relative frequency) for subset of trauma patients
7. *Cumulative* success rates for consecutive ETI attempts
8. Frequencies of critical complications
9. Frequencies of rescue airway use
10. Patients receiving no ETI attempts but in whom airway or ventilatory support is required



**NAEMSP AIRWAY MANAGEMENT REPORTING TEMPLATE**

**Patient demographic information:**

Date: \_\_\_/\_\_\_/\_\_\_ Dispatch Time: \_\_\_:\_\_\_ am / pm  
 EMS Service Name/No: \_\_\_\_\_  
 Pt age (yr): \_\_\_\_\_ Patient sex:  M  F

**1. Indication for invasive airway management (check one):**

- Apnea or agonal respirations
- Airway reflex compromised
- Ventilatory effort compromised
- Injury/illness involving airway
- Adequate airway reflexes/vent effort, but potential for compromise
- Other: \_\_\_\_\_

**2. Was endotracheal intubation (ETI) attempted?**

Yes  No

**3. If ETI not attempted – alternate method of airway support:**

- Bag-Valve-Mask (BVM)  Combitube
- Needle Jet Ventilation  LMA
- Open Cricothyrotomy  Other Cricothyrotomy
- CPAP/BiPAP  Not Applicable (ETI Attempted)
- Other: \_\_\_\_\_

**4-6. Patient subsets (Select Yes/No):**

- Is patient in cardiopulmonary arrest on intubation?  Yes  No
- Is patient a victim of trauma?  Yes  No
- Is patient under 18 years old?  Yes  No

**7-11. Vital signs prior to ETI attempt (leave blank if not obtained):**

Pulse: \_\_\_ beats/min Blood Pressure: \_\_\_/\_\_\_ mmHg  
 Resp Rate: \_\_\_ breaths/min SaO<sub>2</sub>: \_\_\_ %

**12-14. Glasgow Coma Score (GCS) before intubation:**

- Eye:**  none (1)  pain (2)  verbal (3)  spontaneous (4)  
**Verbal:**  none (1)  incomprehensible (2)  
 inappropriate words (3)  
 disoriented (4)  oriented (5)  
**Motor:**  no response (1)  extends to pain (2)  
 flexes to pain (3)  withdraws from pain (4)  
 localizes pain (5)  obeys commands (6)

**15. Monitoring and treatment modalities concurrent with intubation (check all that apply):**

- ECG monitor  Pulse-Oximetry
- IV access  C-spine immobilization
- CPR (chest compressions)  Gum Elastic Bougie
- BAAM  Endotrol Tube
- Other: \_\_\_\_\_

**17. Level of training of each rescuer attempting intubation:**

Rescuer	Level of Training (check one)
A*	<input type="checkbox"/> EMT-P <input type="checkbox"/> EMT-1 <input type="checkbox"/> EMT-3 <input type="checkbox"/> Medic Student <input type="checkbox"/> Nurse/PHRN <input type="checkbox"/> Phys Asst <input type="checkbox"/> MD/DO (attend) <input type="checkbox"/> MD/DO (ms) <input type="checkbox"/> Other: _____
B*	<input type="checkbox"/> EMT-P <input type="checkbox"/> EMT-1 <input type="checkbox"/> EMT-3 <input type="checkbox"/> Medic Student <input type="checkbox"/> Nurse/PHRN <input type="checkbox"/> Phys Asst <input type="checkbox"/> MD/DO (attend) <input type="checkbox"/> MD/DO (ms) <input type="checkbox"/> Other: _____
C*	<input type="checkbox"/> EMT-P <input type="checkbox"/> EMT-1 <input type="checkbox"/> EMT-3 <input type="checkbox"/> Medic Student <input type="checkbox"/> Nurse/PHRN <input type="checkbox"/> Phys Asst <input type="checkbox"/> MD/DO (attend) <input type="checkbox"/> MD/DO (ms) <input type="checkbox"/> Other: _____

**16-18. Provide information for each laryngoscopy attempt. FOR ORAL ROUTE, EACH INSERTION OF BLADE (LARYNGOSCOPY) IS ONE "ATTEMPT." FOR NASAL ROUTE, EACH PASS OF TUBE PAST NARES IS ONE "ATTEMPT."**

Attempt	16. ETI Method	17. Who attempted?	18. Was attempt successful?	Indicate drugs given to facilitate intubation:
#1	<input type="checkbox"/> OTI <input type="checkbox"/> NTI <input type="checkbox"/> Sedation <input type="checkbox"/> RSI <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Midazolam ___mg <input type="checkbox"/> Diazepam ___mg <input type="checkbox"/> Lidocaine ___mg <input type="checkbox"/> Morphine ___mg <input type="checkbox"/> Etomidate ___mg <input type="checkbox"/> Succinylcholine ___mg <input type="checkbox"/> Atropine ___mg <input type="checkbox"/> Topical Spray <input type="checkbox"/> Other - Specify: _____ mg <input type="checkbox"/> Other - Specify: _____ mg
#2	<input type="checkbox"/> OTI <input type="checkbox"/> NTI <input type="checkbox"/> Sedation <input type="checkbox"/> RSI <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No	
#3	<input type="checkbox"/> OTI <input type="checkbox"/> NTI <input type="checkbox"/> Sedation <input type="checkbox"/> RSI <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No	
#4	<input type="checkbox"/> OTI <input type="checkbox"/> NTI <input type="checkbox"/> Sedation <input type="checkbox"/> RSI <input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No	

**19-24. Endotracheal tube confirmation:**

- 19. Auscultation  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- 20. Bulb Aspiration  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- 21. Syringe Aspiration  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- 22. Colorimetric ETCO<sub>2</sub>  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- 23. Digital ETCO<sub>2</sub>  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- 24. Waveform ETCO<sub>2</sub>  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.
- Other: \_\_\_\_\_  Tracheal Placement  Esophageal Placement  Indeterminate  Not Assessed  Tube not placed.

25. Peak ETCO<sub>2</sub> value: \_\_\_\_\_  Indeterminate

**26. Was ETI successful for the overall encounter (on transfer of care to ED or helicopter)?**

Yes  No

**27. Who determined the final placement (location) of ET tube?**

- Rescuer performing intubation.
- Another rescuer on the same team.
- Receiving helicopter crew.
- Receiving hospital team.
- Other: \_\_\_\_\_

**28-32. Vital signs after intubation attempt:**

Pulse: \_\_\_ beats/min Blood Pressure: \_\_\_/\_\_\_ mmHg  
 Resp Rate: \_\_\_ breaths/min SaO<sub>2</sub>: \_\_\_ %

**33. Critical complications encountered during airway management (Check all that apply):**

- Failed intubation effort.
- Injury or trauma to patient from airway management effort.
- Adverse event from facilitating drugs.
- Esophageal intubation - delayed detection (after tube secured)
- Esophageal intubation - detected in ED.
- Tube dislodged during transport/patient care.
- Other: \_\_\_\_\_

**34. If all intubation attempts FAILED, indicate suspected reasons for failed intubation (check all that apply):**

- Inadequate patient relaxation  Orofacial trauma.
- Inability to expose vocal cords  Secretions/blood/vomit.
- Difficult pt anatomy.  Unable to access pt.
- ETI attempted, but arrived at destination facility before accomplished.
- Not applicable - Successful field ETI  Other: \_\_\_\_\_

**35. If all intubation attempts FAILED, indicate secondary (rescue) airway technique used (check all that apply):**

- Bag-Valve-Mask (BVM) Ventilation  Needle/Jet Ventilation
- Combitube  Open Cricothyrotomy
- Not applicable - Successful field ETI  Other: \_\_\_\_\_

**36. Did secondary (rescue) airway result in satisfactory ventilation?**

Yes  No  Not applicable

**37-38. Airway Management Times**

Time of decision to intubate: \_\_\_:\_\_\_ am / pm  
 Time of successful intubation: \_\_\_:\_\_\_ am / pm  
 Time intubation abandoned: \_\_\_:\_\_\_ am / pm

Wang, H. E., A. B. Peitzman, et al. (2004). "Out-of-hospital endotracheal intubation and outcome after traumatic brain injury.[see comment]." Annals of Emergency Medicine 44(5): 439-50.

(n=2,301, 56.1%). Adjusted odds of death were higher for out-of-hospital endotracheal intubation than ED endotracheal intubation (odds ratio [OR] 3.99; 95% confidence interval [CI] 3.21 to 4.93). Out of hospital endotracheal intubation was associated with an increased adjusted odds of poor neurologic outcome (OR 1.61; 95% CI 1.15 to 2.26), moderate or severe functional impairment (Functional Impairment Score 6 to 15; OR 1.92; 95% CI 1.40 to 2.64), and severe functional impairment (Functional Impairment Score 11 to 15; OR 1.80; 95% CI 1.29 to 2.52). CONCLUSION: Out-of-hospital endotracheal intubation was associated with adverse outcomes after severe traumatic brain injury. The implications for current clinical care remain undefined.

Wang, H. E., D. F. Kupas, et al. (2003). "Preliminary experience with a prospective, multi-centered evaluation of out-of-hospital endotracheal intubation." Resuscitation 58(1): 49-58.

median per-service return rate was 75%. Non-response (data form not returned for attempted intubation) was problematic, with nine services demonstrating data return rates less than 50%. Data return rates could not be calculated for an additional nine services. The missing data entry rate was 0.5-22.2%. The overall reported ETI success rate was 86.8% (92.8% for cardiac arrests and 76.8% for non-arrests) and did not appear to vary between population settings. There were two cases of delayed

Wayne, M. A. and E. Friedland (1999). "Prehospital use of succinylcholine: a 20-year review.[see comment]." Prehospital Emergency Care 3(2): 107-9.

**RESULTS:** Paramedics successfully intubated 95.5% (1,582) of all patients receiving succinylcholine, 94% (1,045) of trauma patients, and 98% (538) of medical patients. They were unable to intubate 4.5% (74) of the patients. All of these were successfully managed by alternative methods. Unrecognized esophageal intubation occurred in six (0.3%) patients. The addition of capnography and a tube aspiration device, in 1990, decreased the incidence of esophageal intubations.

**CONCLUSION:** Paramedics trained to use succinylcholine, to assist the process of endotracheal intubation, can safely intubate a high percentage of patients.

**What does  
all of this  
mean?**



**Endotracheal intubation  
in the field may be  
rarely indicated**

**Endotracheal intubation  
in the field may be  
harmful...**

**...indeed, if you tracked your  
own survival data,  
you would likely find that  
people were less likely to survive  
with Field ETI**

**Why?**

# Field ETI

- **Prolonged attempts**
- **Hypoxia during attempts**
- **Multiple attempts**
- **Aspiration during attempts**
- **Hyperventilation AFTER intubation**
- **Instrumenting the airway  
in critical patients**

# **Example**

**Medic reports to Medical Director that an elderly patient in respiratory distress was cared for in the field, given nebs and oxygen, improved to GCS 8**

**On arrival to ED, Doc takes one look at her and makes 8 attempts to intubate her, during which she bradys down and dies**



# Example

**What does that mean?**

**Did that patient need to be intubated who was improving with oxygen and supportive care?**

# **Fowler's Maxim**

**“The First Five Minutes”**

# Fowler's Maxim

In some patients you have  
to act right now:

*Airway obstruction*

*Exsanguination*

*Cardiac Arrest*

*Profound respiratory distress*

*Shock*

# Fowler's Maxim

**In most patients, you have  
five minutes**

**The patient who seems stable  
for the moment, with a decent pressure,  
with a decent pulse ox, whose airway  
is not immediately threatened**

**TAKE A MINUTE OR TWO TO THINK**

# **Delaney's Corollary**

**If it took them a couple of days or more to get sick, then you probably have at least five minutes to stop and think**

**While you're getting oxygen, IV's, other supportive care, arranging for transport to the appropriate facility**





**The Airway of the future  
will be fast, effective,  
and virtually hazard free**

- **Avoiding airway trauma**
- **Virtually always goes in**
  - **Easy to train**
  - **Easy to remember**



# The Easy Tube



# The Easy Tube



**The Airway of the future  
will be fast, effective,  
and virtually hazard free**



# Breathing and Circulation

# **Resuscitation Outcomes Consortium**

**A consortium of ten cities and states  
across North America  
designated by the National Institutes of Health  
to be the largest EMS research group  
in the history of medicine**

# **Resuscitation Outcomes Consortium**

**Will be initially looking at hypertonic saline infusions for traumatic brain injury and hemorrhagic shock due to trauma**

**Already in use in the military, just not studied in large civilian populations**



# **Resuscitation Outcomes Consortium**

**Will next look at the  
Impedance Threshold Device  
to improve negative intrathoracic pressure  
during cardiac compressions**

# **Resuscitation Outcomes Consortium**

**Impedance Threshold Device**

**Turns chest compressions from a  
“one stroke engine” to a  
“two stroke engine”**

# **Resuscitation Outcomes Consortium**

## **Impedance Threshold Device**

**Makes use of chest recoil with  
tiny, momentary airway occlusion  
to increase venous return**

# **Resuscitation Outcomes Consortium**

## **Impedance Threshold Device**

**Essentially normalizes blood pressure  
during chest compressions**

## Impedance Threshold Device May Improve Survival in Out-of-Hospital Cardiac Arrest

**Peggy Peck**

Nov. 12, 2004 (New Orleans) — Use of an investigational inspiratory impedance threshold device (ITD) — ResQPod Circulatory Enhancer — during standard cardiopulmonary resuscitation (CPR) was associated with a doubling of short-term survival in patients with pulseless electrical activity (PEA) at any time during resuscitation, according to study results presented here at the American Heart Association 2004 Scientific Session.

Lead investigator Tom P. Aufderheide, a professor of emergency medicine at the Medical College of Wisconsin in Milwaukee, told Medscape he predicts the device "will have as great an impact on [CPR] efforts and treatment of out-of-hospital cardiac arrest as AEDs." He said use of the device increases blood supply to the brain and heart during the resuscitation.

The study compared the device, made by Advance Circulatory Systems Inc. in Eden Prairie, Minnesota, to a sham device for use during standard manual CPR. The device can be used with a face mask or can be attached to an endotracheal tube, and it is equipped with a timing light that flashes at a rate of 12 breaths a minute with each breath lasting a maximum of 1.5 seconds.

The primary end point was admission to an intensive care unit.

There were 116 patients randomized to the sham device and 114 were randomized to the ITD. Of the patients randomized to the active device, the one-hour survival rate was 26%, the ICU admission rate was 25%, and 24-hour survival rate was 17%. For patients in the sham device group the rates were lower — 18%, 17%, and 12%, respectively — but the differences were not statistically significant.

### Information from Industry

**[St. Jude Medical Advanced Solutions for Cardiac Rhythm Disorders](#)** - Get the latest on Heart Failure and Atrial Fibrillation therapies and devices, technical insights, patient information, and more.

**[Cialis® \(tadalafil\)](#)**

**[ULTRACET \(37.5 mg tramadol HCl / 325 mg acetaminophen tablets\) for short-term management of acute pain](#)**

[Other Product InfoSites](#)

## ResQPOD Circulatory Enhancer<sup>®</sup>

### Introducing the Most Advanced Device for Enhancing Circulation during Cardiopulmonary Resuscitation (CPR)

The ResQPOD Circulatory Enhancer\*

- Increases blood flow to the heart and brain during CPR
- Increases the opportunity for survival and normal neurological outcome
- May be used with standard CPR or pump-assisted (active compression decompression - ACD) CPR
- Works in conjunction with standard resuscitation techniques and equipment





# Physiology

**Oxygen** -> lungs -> alveoli -> blood

breath  
↑  
**CO<sub>2</sub>**

lungs

↑  
**CO<sub>2</sub>**

blood

blood

**Oxygen**

↓  
muscles + organs

**Oxygen**


↓  
cells

**Oxygen**  
+  
**Glucose**

energy

←  
**CO<sub>2</sub>**

3/29/2006



The negative pressure inside the thorax “pulls” blood back from the positive pressure areas.

Positive pressure

Negative pressure

Positive pressure



Maintaining the “negativity” of the pressure inside of the thorax is one of the most vital areas of understanding resuscitation

Negative pressure

3/29/2006






# Signs of Shock

**Early**



**Weak, thirsty, lightheaded**  
**Pale, then sweaty**  
**Tachycardia**  
**Tachypnea**  
**Diminished urinary output**

**Late**



**Hypotension**  
**Altered LOC**  
**Cardiac arrest**  
**Death**


# What does a low blood pressure mean?

*Either...*

*Or a combination  
of any of these*

*...from BTLS, editions 2, 3, 4, and 5 Fowler et al*

# Shock




## Cardiogenic

Rapid pulse  
Distended neck veins  
Cyanosis



## Volume Loss

Rapid pulse  
Flat neck veins  
Pale



## Vasodilatory

Variable pulse  
Flat neck veins  
Pale or pink

**We have been overventilating  
patients in circulatory collapse  
to death for years, without  
knowing it**

*“After all, if a little oxygen  
is **GOOD**,  
more is better, right?”*



**WRONG!**

**Overventilation raises  
intrathoracic pressure,  
decreasing venous return,  
and dropping cardiac output**

Blood pressure =

(Cardiac output) x  
(Volume) x  
(Peripheral resistance)

**The heart only pumps out  
what it gets back**

**If you drop venous return,  
cardiac output drops**

**That is, if the pump can't fill,  
then the pump can't pump**

# Detrimental hemodynamic effects of assisted ventilation in hemorrhagic states

Paul E. Pepe, MD, MPH; Keith G. Lurie, MD; Jane G. Wigginton, MD; Claus Raedler, MD; Ahamed H. Idris, MD

**Objective:** Our goal was to demonstrate explicitly that lower-frequency positive-pressure ventilation not only preserves adequate oxygenation and acid-base status in hemorrhagic states, but also that “normal” or higher respiratory rates significantly compromise hemodynamics, even with moderate degrees of hemorrhage.

**Design and Subjects:** Eight intubated pigs (ventilated with 12 mL/kg tidal volume, 28% F<sub>IO</sub><sub>2</sub>, respiratory rate = 12 breaths/min) were hemorrhaged to <65 mm Hg of systolic blood pressure. Respiratory rates were then sequentially changed every 10 mins to 6, 20, 30, and 6 breaths/min.

**Measurements and Main Results:** With respiratory rates at 6 breaths/min, all subjects maintained pH of >7.25 and Sa<sub>O</sub><sub>2</sub> of >99% while increasing systolic blood pressure (mean, 65–84 mm Hg;  $p < .05$ ), time-averaged coronary perfusion pressure ( $50 \pm 2$  to  $60 \pm 4$  mm Hg;  $p < .05$ ), and cardiac output (2.4 to 2.8 L/min;  $p < .05$ ). With respiratory rates of 20 and 30 breaths/min, systolic blood pressure ( $73 \pm 4$  and  $66 \pm 5$  mm Hg, respectively),

coronary perfusion pressure ( $47 \pm 3$  and  $42 \pm 4$  mm Hg), and cardiac output (2.5 and 2.4 L/min) diminished. When returned to 6 breaths/min, systolic blood pressure (95 mm Hg), coronary perfusion pressure ( $71 \pm 6$  mm Hg), and cardiac output (3.0 L/min) improved significantly ( $p < .05$  for all comparisons).

**Conclusions:** After moderate hemorrhage, animals maintain adequate oxygenation and acid-base status with lower-frequency respiratory rates, whereas increasingly higher respiratory rates progressively and significantly impair hemodynamics. Current ventilatory protocols for trauma resuscitation should be re-examined and considered a possible cause of worsened clinical outcomes and unrecognized confounded study results. (Crit Care Med 2004; 32[Suppl.]:S414–S420)

**KEY WORDS:** hemorrhage; shock; hypovolemia; hemorrhagic shock; ventilation; positive-pressure ventilation; hemodynamics; mechanical ventilation; auto-positive end-expiratory pressure; preload; venous return; coronary perfusion pressure; respiratory support

Crit Care Med 2004 Vol. 32, No. 9 (Suppl.)






Figure 2. Digital recording of (A) effect of 12 positive-pressure breaths/min on aortic pressures in swine after a moderate degree of hemorrhage, demonstrating intermittent depressions in these pressures immediately after each breath (refer back to Fig. 1). Breaths are indicated in the *upper tracing* (airway pressure curve). B, effect of changing to 6 positive-pressure breaths/min, demonstrating less intermittent depression and relatively higher levels of aortic pressures when compared with baseline respiratory rates of 12 breaths/min (A). C, effect of changing from 6 to 30 positive-pressure breaths/min, showing return of significant depressions in aortic pressures that are much lower than those observed with respiratory rates of 6 and 12 breaths/min (A and B). D, effect of returning to 6 positive-pressure breaths/min, showing dramatic elevations in aortic pressures and substantial enhancements of the corresponding area under the aortic diastolic pressure curve, reflecting marked improvements in summary coronary perfusion pressures.



**Research is clear on the fact  
that all emergency providers)  
bag patients too fast  
with too BIG of a squeeze  
and too FAST on the squeeze**

**Slow down the rate of ventilation  
until capnography begins to rise**

**Maintain a minute ventilation  
of approximately five liters and  
see where capnography goes  
from there**

**One hand squeeze every 8 seconds**

# CPAP

**Vitaid** *Aiding Life* 

Home | Products ▾ | Departments ▾ | Order | About Us | Contact Us | Search

**Boussignac CPAP Resource Center**

- Boussignac CPAP System
- Boussignac CPAP + Nebulizer
- FAQ
- Education

**Boussignac CPAP System**

**For Acute Pulmonary Edema (APE)**

**Lowest O2 consumption rate of any CPAP system**

- Only 15-30L/min flow rate for 3.5-10cmH2O

**System doesn't require a flow generator**

- Eliminates investment in capital equipment and repair
- Boussignac CPAP only needs O2 source and flowmeter

**Only completely open CPAP system**


- Eliminates re-breathing
- Reduces risk of barotrauma
- Decreases the work of breathing
- Accommodates patient's high peak inspiratory flow demand
- Allows use of suction catheter or bronchoscope without loss of CPAP

**Only CPAP system that depends entirely on the O2 flow rate**

- CPAP easily adjusts to patient's need (2.5-20cmH2O) simply by titrating O2 flow
- PEEP adjustable by changing flow rate, no valves to change
- Constant, accurate pressure measurement via optional manometer

**Complete, easy to use, portable, single use system**

- CPAP set up in <2 minutes, therefore ideal for all EMS and ED settings
- Requires 75% less space in field packs than competitive systems
- Boussignac CPAP weighs only 10grams, 2" long
- Permits CPAP without interruption during transport



## Boussignac CPAP works the same way as the turbines of a jet engine.


**OXYGEN SUPPLY**  
Oxygen molecules enter the chamber

**OXYGEN ACCELERATION**  
The molecules of oxygen are accelerated at the speed of sound as they pass through micro channels




**OXYGEN BRAKING**  
The molecules of oxygen strike a deflector which sends them back to the central zone (mixing zone)


**ESTABLISHMENT OF A VIRTUAL VALVE**  
The collision of the molecules generates a turbulence which transforms the speed into pressure





# What about the AutoPulse?







# Drugs and ECG's

# Drugs

*It is clear that overventilation  
has changed patient outcomes  
over the years and  
prevented us from seeing what  
drugs could*

# **Which Drugs to Revisit?**

**Antiarrhythmics in arrest**

**Pressors during arrest**

# **Future Drugs**

**Artificial hemoglobins?**

**Vasopressin?**



*...but just when we thought  
life was getting easier...*

# **Electrocardiography**

**12 Lead Interpretation**

**by medics**

**is now the standard of care**

**even if the 12 lead is NOT**

**on your rig**

# ECG Interpretation

The image features a central orange square with a dark blue border. Inside the orange square, there is a faint, glowing yellow anatomical diagram of the heart and its major vessels. Overlaid on this diagram is a black ECG waveform with a red line tracing its path. The text 'ECG Interpretation' is written in a bold, yellow, sans-serif font at the top. Below it, the phrase 'Anatomically speaking...' is written in a white, italicized, serif font, slanted diagonally across the center.

*Anatomically  
speaking...*





**SA**

**Bundle of His**

**AV**

**Bundle  
Branches**





Name:	CLEMENTS, JULIA	12-Lead 1	HR 169bpm	Abnormal ECG <b>**Unconfirmed**</b>
ID:	101406124512	10/14/2006	12:48:15 PM	wide QRS tachycardia
Patient ID:		PR 0.000s	QRS 0.188s	Left bundle branch block
Incident ID:	615292	QT/QTc:	0.350s/0.586s	
Age: 65	Sex: F	P-QRS-T Axes:	0 0 118	




x1.0 05-150Hz 25mm/sec  
Medtronic, Inc. Comments:

A-9 000 3011371-124 LP1231254529

File Edit View Help

Name:	CLEMENTS, JULIA	12-Lead 1	HR 169bpm	Abnormal ECG <b>**Unconfirmed**</b>
ID:	101406124512	10/14/2006	12:48:15 PM	wide QRS tachycardia
Patient ID:		PR 0.000s	QRS 0.188s	Left bundle branch block
Incident ID:	615292	QT/QTc:	0.350s/0.586s	
Age: 65	Sex: F	P-QRS-T Axes:	0 0 118	



Device	Device ID	New	Report	*	Ti...	Time	Elapsed Time	Event Type	*	Note	HR	SpO2*PR	EtCO2(mmHg)*RR	NIBP(mmHg)*PR	P1(mmHg)
LP12	A-9	⊗	Continuous Complete	⊗	0...	12:45:21	00:00:00	Low Battery	♥						
LP12	A-9	⊗	Start Trend Data		0...	12:45:21	00:00:00	Power On	♥						
LP12	A-9	⊗	Initial Rhythm		0...	12:45:21	00:00:00	Start Trend Data	♥						
LP12	A-9	⊗	12-Lead 1		0...	12:45:52	00:00:31	Initial Rhythm	♥						
						12:48:15	00:02:54	12-Lead 1	♥						
						12:50:19	00:04:58	Vital Signs	♥		167				
						12:55:19	00:09:58	Vital Signs	♥		167				
						13:00:19	00:14:58	Vital Signs	♥		169				
						13:00:26	00:15:05	Power Off	♥						



# 66.34.180.153 - Remote Desktop

Security Configurati...

NotePad

My Document...

DALLAS\_20...

My Comput...

My Netwo...

Internet Explorer

d2d.web.rar

CMD

d2d.web.rar

D:\INETPUB\UTSOUTHWESTERN\FTP\archives\30200\Garland\30200.102006

File Edit View Favorites Tools Help

Back Search Folders Go

Address D:\INETPUB\UTSOUTHWESTERN\FTP\archives\30200\Garland\30200.102006

Folders	Name	Size	Type	Date Modified
deploy	30200.2006102720165100LP...	360 KB	PCO file	10/28/2006 4:10
ire	30200.2006102707262300LP...	31 KB	PCO file	10/28/2006 4:10
lib	30200.615931_615931.pco	327 KB	PCO file	10/27/2006 4:10
logs	30200.2006102417533700LP...	319 KB	PCO file	10/27/2006 4:10
security	30200.449213059_20061024...	429 KB	PCO file	10/27/2006 4:10
Temp	30200.2006102420353500LP...	230 KB	PCO file	10/26/2006 4:10
Tools	30200.2006102418300800LP...	434 KB	PCO file	10/26/2006 4:10
INETPUB	30200.2006102204102300LP...	142 KB	PCO file	10/26/2006 4:10
UTSOUTHWESTERN	30200.2006102204102300LP...	142 KB	PCO file	10/26/2006 4:10
FTP	30200.2006102203590800LP...	34 KB	PCO file	10/26/2006 4:10
archives	30200.2006102115090700LP...	31 KB	PCO file	10/26/2006 4:10
30000	30200.2006101519311200LP...	31 KB	PCO file	10/26/2006 4:10
30100	30200.2006101507540500LP...	30 KB	PCO file	10/26/2006 4:10
30200	30200.2006101422424200LP...	179 KB	PCO file	10/26/2006 4:10
Garland	30200.450640315_615831.pco	210 KB	PCO file	10/26/2006 4:10
30200.032006	30200.2006102412391000LP...	392 KB	PCO file	10/25/2006 4:10
30200.052006	30200.2006101516390500LP...	287 KB	PCO file	10/25/2006 4:10
30200.062006	30200.2006100913394400LP...	14 KB	PCO file	10/25/2006 4:10
30200.072006	30200.2006100913394400LP...	366 KB	PCO file	10/25/2006 4:10
30200.082006	30200.449688909_615117.pco	41 KB	PCO file	10/25/2006 4:10
30200.092006	30200.444545331_615127.pco	286 KB	PCO file	10/25/2006 4:10
30200.102006	30200.412468472_615212.pco	37 KB	PCO file	10/25/2006 4:10
30300	30200.172162383_615219.pco	111 KB	PCO file	10/25/2006 4:10
DeSoto	30200.2006102316385500LP...	308 KB	PCO file	10/24/2006 4:10
30300.102006	30200.2006102120412000LP...	393 KB	PCO file	10/23/2006 4:10
30400	30200.2006102107421900LP...	244 KB	PCO file	10/23/2006 4:10
30500	30200.2006101707112000LP...	38 KB	PCO file	10/23/2006 4:10
30600	30200.2006101504170000LP...	252 KB	PCO file	10/23/2006 4:10

Unconfirmed\*\*

cardia

h block

Note HR SpO2\*PR

Shortcuts

Windows Update

Help and Support

Mouse

Network Connections

Phone and Modem ...


Portable Media Devices

Power Options

Printers and Faxes


Regional and Language ...

# Multifocal Atrial Tachycardia





# IMMEDIATE Trial



# **IMMEDIATE Trial**

**Medic identifies the Acute Coronary Syndrome through patient History and Physical and through 12 Lead Interpretation**

**Medic gets consent**

**Medic initiates an infusion of Glucose, Insulin, and Potassium in the field to limit incidence of sudden death and infarct size**



# **IMMEDIATE Trial**

**Over time EMS will likely be giving medication to limit infarct size and incidence of sudden death**

**This will ultimately increase survival from acute coronary syndrome and decrease the burden of CHF post infarct to patients and the community**

# **IMMEDIATE Trial**

**Turns out that sick tissue  
likes a little extra insulin around**

**Insulin turns off fat metabolism and  
turns on carbohydrate metabolism,  
giving more energy per gram of substrate  
to sick cells**

**This has been know for over 40 years**

# **IMMEDIATE Trial**

**Is that cool,  
or what???**


**Speaking of  
Cool?**

**How 'bout  
some cases?**




**What is the patient's  
blood pressure?**

**A 15 year old AA male  
is found confused, sweaty, with  
a respiratory rate of 36,  
a systolic pressure of 80, and  
this EKG rhythm strip**






**60 year old Aunt Minnie  
presents with systolic of 90  
and no cardiac history**



**She has been ill for two days**

**60 year old with rate of 158**


$$220 - 60 = 160$$



**What statement can you make?**

**60 year old with rate of 158**


$$220 - 60 = 160$$



**Does she need Adenosine?**

# Synthesis






So,  
Who's  
Foolin'  
Who??



**Emergency providers  
are primary members  
of the medical team.**

**The scope of practice  
of these professionals  
continues to grow  
with passing years**





**Let us then**  
apply our best efforts  
to monitor emerging research  
with the sharpened focus  
of clarity and simplification,  
pooling our individual creativities  
for the greater good  
of those we serve.





**This Talk may be found at**

**[www.rayfowler.com](http://www.rayfowler.com)**

***[drray@doctorfowler.com](mailto:drray@doctorfowler.com)***

... *and Good Afternoon!*

*Questions or comments?*



